

## **Welfare and Conservation Implications of Intentional Breeding for the Expression of Rare Recessive Alleles**

Association of Zoos & Aquariums

Animal Welfare Committee: Taskforce on Animal Breeding Practices

Approved by the AZA Board of Directors – June 2011

### **AZA Position Summary:**

Breeding practices that increase the physical expression of single rare alleles (i.e., rare genetic traits) through intentional inbreeding, for example intentional breeding to achieve rare color-morphs such as white tigers, deer, and alligators, has been clearly linked with various abnormal, debilitating, and, at times, lethal, external and internal conditions and characteristics, which are outlined in this paper. Many of these conditions may seriously compromise the welfare of individual animals. In addition, such breeding practices are also problematic from a population management and conservation perspective, impairing our ability to develop and maintain sustainable captive populations for the future and to deliver appropriate animal welfare and conservation education messages.

Therefore such practices are not in adherence with AZA's Board-approved Policy on the Presentation of Animals (22 July 2008), which maintains that "... animals should always be presented according to the following core principles: 1. Animal and human health, safety, and welfare are never compromised; 2. Education and a meaningful conservation messages are integral components of the presentation; and, 3. The individual animals involved are consistently maintained in a manner that meets their social, physical, behavioral, and nutritional needs."

Based on the welfare, population health and management, as well as conservation and education concerns outlined in this paper, AZA-accredited institutions should not engage in intentional breeding practices for the purpose of producing anomalous phenotypes.

At times, inappropriate breeding practices by others (outside of AZA) may yield animals with anomalous phenotypes and adverse internal conditions, which may be in need of rescue. Providing holding and care for such animals and responding to rescue requests from local, state, or federal agencies are appropriate activities for AZA-accredited institutions, provided that the delivery of thoughtful educational messages about the unfortunate results of intentional inbreeding for rare genetic traits are part of any public display.

---

Interestingly, the very instinct that appears to draw humans towards novel patterns and diversity in general also seems to underlie our fascination with unusual and abnormal patterns and phenotypes expressed only rarely, or occasionally, in nature. The spectacle provided by displays of calves with two heads, five toed cats, and traits such as albinism, melanism, or dwarfism, continues, even today, to provide an attraction to many, unaware of the biology underlying such odd occurrences. Even among today's frequently well informed and educated zoo visitors, the interest in seeing white tigers, white lions, white alligators, or king cheetahs continues often in preference over the 'normal' looking individuals of the same species.

Of greater concern, in some cases, there exists the misconception that these unusual color morphs, or other phenotypic aberrations, may represent a separate endangered species in need of conservation. Various articles and education efforts over the years have tried to put this issue into a proper scientific perspective, at least for the tiger (e.g., Leyhausen & Reed, 1971; Latinen 1987; Schroeter, 1981; EAZA 2010). However, some of the myths persist and will require additional education and a clear stance on this issue from the AZA community.

The purpose of this White Paper, therefore, is to highlight some of the well documented welfare and conservation concerns involved in breeding, or support of breeding, for unusual and rare traits (in any species) and provide recommendation with regard to this practice for AZA accredited institutions and certified related facilities.

A precedent for such recommendations has been set in the area of pet and farm animal welfare in Europe and the UK, where various legislative efforts are under way, or have already been put in place, to discourage the intentional breeding for rare recessive alleles with known potential deleterious effects and welfare concerns. A long list of potentially harmful recessive traits exists for pet animals, from fish and hamsters to dogs and horses, that are associated with either substantial physical or behavioral impairment and/or even lethal outcomes such as a drastically shortened lifespan (e.g., Not et al., 2008; Stucki et al., 2008; Kirkwood et al., 2010). For example, taillessness in some duck breeds can lead to lowered reproductive rates and shorter lifespan as well as problems with copulation and egg laying (Stucki et al., 2008).

Lack of pigmentation in snakes has been associated with detrimental changes in skin and eyes, such as reduced heat exchange ability and neural issues (Not, 1998). Dwarfism in rabbits has been associated with problems in teeth positioning and eating, as well as thermoregulation and reproduction. Indeed most dwarf forms of a variety of species seem to be more prone to general infections and may show compromised immunity (Not et al., 2008). In cats, breeds such as the Manx cat and tailless cat are associated with locomotive disorders, dispositioning of the vertebral column, difficulties defecating, and a loss of about a quarter of offspring when breeding for the trait of "taillessness". The "dominant white" trait in domestic cats (in spite of the word "dominant" in the trait's name the individuals are actually homozygous for this recessive color morph) has been associated with increased occurrence of deafness. Similarly in dogs, the "Merle factor" (e.g. Blue Merle Collie, or Merle Bobtail) has been associated with a disposition to deafness and eye disorders (Steiger et al., 2008). The list is

lengthy and several comprehensive reviews have documented the various problems associated with intentional breeding for rare and recessive alleles in a wide variety of species (e.g., Not et al., 2008; Steiger 2005, 2008; Steiger et al. 2008; Stucki et al., 2008; Kirkwood et al., 2010; Rooney and Sargan, 2010).

As a consequence of these findings, a declaration of intent was adopted in 1995 by the European Convention for the Protection of Pet Animals based on a multilateral consultation with stake holders. Some of the highlights of the declaration included an agreement to take necessary measures to control the breeding of animals that show genetic or phenotypic characteristics harmful to the welfare of the animals in order to prevent suffering, and to develop educational information for the public regarding these issues (website; Steiger et al., 2008).

Similarly, in 1999, the Federation of Veterinarians of Europe (FVE) urged its member countries and the European Commission to consider the introduction of measures designed to safeguard the welfare of animals with respect to the risks inherent in selective breeding for rare and recessive traits. It was stated that this form of selective breeding may cause welfare problems of the following types: offspring produced may be unable to express their natural behaviors and/or may be predisposed to a variety of hereditary, congenital, metabolic or infectious disease, disability, and early death (FVE, 1999; Steiger et al. 2008). Since then various European countries have indeed implemented legislative standards for animal breeding and have even gone so far as to outlaw breeding for some of the more extreme traits, such as intentional breeding of Manx cats or 'Merle factors' in dogs (e.g., Germany, 2000; Austria, 2005).

In exotic species hereditary problems associated with selective breeding have also been clearly documented. For example, most white tigers currently in captivity are Amur-Indian hybrids that have been highly inbred to achieve continued occurrence of the colormorph (Thornton et al., 1967; Thornton, 1978; Roychoudhury and Sankhala, 1979). Various abnormalities and deformities associated with such selective breeding practices have been documented, such as the occurrence of an abnormality of the visual pathways in the brain, resulting in visual impairments such as strabismus, a condition that involves a lack of proper alignment of the eyes, preventing binocular focus on any particular point or object, and thus negatively affecting depth perception (Guillery and Kaas, 1973). This congenital defect has been listed as a common abnormality also found in Siamese cats, and in albino ferrets, albino mink, and other albinos of various mammal species that have been studied (Creel and Giolli, 1972; Sanderson and Guillery, 1973; Guillery and Kaas, 1973; Guillery, 1986).

Vascular ring anomaly around the trachea and esophagus has also been reported. This abnormality leads to the inability to feed and swallow effectively and requires an operation for correction and survival (Ketz et. al. 2001). This type of abnormality similarly represents a common congenital problem in domestic dogs and cats (e.g., Fox, 1988) resulting from inbreeding. The same abnormality has also been reported in white lions (Goldin and Lambrechts, 1999). Other congenital defects, such as changes in cranial structures and skull development, as well as cleft palate have also been well documented in white tigers and other rare color morphs in other species. More generally,

albinism has been associated with a wide variety of health problems and congenital defects and is regarded as a hereditary defect rather than a desirable trait in wild and most domestic populations (Creel and Giolli, 1972; Guillery, 1986; Laikre, 1999).

The underlying cause for the multitude of the above cited health and welfare issues is a relatively simple one. Several of the traits such as albinism are located on recessive alleles that are only expressed phenotypically (in appearance) if two copies of the same trait are obtained by a given offspring (homozygous representation of alleles). For example, in the case of the white tiger one recessive allele has to come from each parent to allow for expression of the white striped color morph. While this has happened rarely in wild tiger populations, such as one in India many decades ago, and may occur in wild populations of various species occasionally (e.g., white deer, lion, ferret), such traits only rarely get expressed, and, when expressed, it is very likely that they confer a disadvantage resulting in reduced fitness for a given individual under most circumstances. Indeed, the very rarity of the traits in natural populations is itself strong evidence that they have deleterious consequences.

A wide variety of heritable defects and abnormalities have been found associated with recessive alleles, and the occurrence of these detrimental and/or lethal recessive traits is termed the genetic load of a population and species. Since these alleles are generally rare in a given population, some level of inbreeding (i.e., breeding of closely related individuals, which increases the chance that the same two rare and recessive alleles will be present in the two parents and therefore can be transmitted to and expressed in offspring) has to be practiced to achieve phenotypic expression of the trait with greater than a miniscule frequency of occurrence. Selective breeding for such traits will therefore usually lead to a variety of negative consequences also documented for severe inbreeding, such as increased expression of the genetic load, congenital defects, such as cleft palate, and overall decline in fecundity and increase in morbidity (e.g., Falconer, 1981). In summary, the welfare concerns associated with intentional breeding for rare (and thus usually detrimental) traits are therefore twofold:

1. Health and welfare problems directly associated with the trait itself, such as visual and neural problems associated with albinism, or gait and elimination problems associated with taillessness.
2. The sometimes more indirect, but just as problematic, health and welfare consequences related to intensive inbreeding to accomplish expression of rare and unusual traits, such as congenital deformities, decline in overall fitness and fertility, increased susceptibility to disease and infection and shortened lifespan or still birth.

Furthermore, in terms of effective conservation management and population health, selective breeding for specific phenotypes is in direct opposition to standard zoo population management goals of maximizing genetic diversity by minimizing inbreeding. Recessive alleles (or any particular alleles) should neither be selected against nor selected for, since doing so would lead to a loss of overall genetic diversity (Lacy, 2000).

Selection for specific traits will hasten a population's loss of gene diversity, lead to higher inbreeding levels more quickly, and create a domesticated form of the species that no longer represents or resembles the wild population.

Occasional expression of a rare and deleterious allele may occur by chance, but should not be 'forced'; rather, these traits should be allowed to appear at their naturally occurring frequencies. Retaining all alleles at their naturally occurring frequencies helps retain genetic diversity and provides populations with adaptive potential in the face of environmental change. The standard genetic management strategy in zoos, using mean kinship rather than phenotype to select breeding animals, is specifically intended to maintain gene diversity and minimize inbreeding (Ballou & Lacy, 1995). The mean kinship genetic management strategy aims to efficiently equalize all founder lineages in small populations, whereas preferential breeding of rare or abnormal phenotypes within the limited space in zoos causes over-production of a few lineages at the expense of known-pedigreed animals from genetically valuable lineages. Inequality of founder lineages will lower genetic diversity at the population level and will eventually lead to individual genetic problems associated with inbreeding.

Modern zoos are concerned with future sustainability of wildlife populations. Problems for many captive populations arise from the small genetic base (i.e., few founder individuals) from which zoo breeding programs often are started, the difficulty of obtaining breeding activity reliably and successfully, and the small populations that can be maintained with the resources available for most programs. It is especially counter-productive therefore, to allocate breeding program resources toward practices that rapidly degrade the genetic variation with which the program was started, cause reduced reproductive fitness, shorter life spans, and reduce the resources that can be allocated for animal management programs that do serve bona fide conservation and education purposes.

From an education perspective, intentional inbreeding for the production of anomalous phenotypes is in direct contrast to the mission of modern zoos. Propagating animals that specifically do not represent the normal characteristics and variation of the species creates a confused educational message. If animals in zoos and aquariums are to be ambassadors for their species then the exhibition of phenotypic aberrations creates difficulty in properly interpreting what modern zoos are about. Further, it is counter-productive to the overall conservation message of preserving essential characteristics of a species and ensuring genetically healthy and sustainable populations both in zoos and aquariums and in the wild.

Primary roles of modern zoos are to help protect wildlife and natural habitat, educate the public about conservation efforts, and directly engage the public in these efforts. These commendable goals are forfeited when intentional inbreeding is practiced to create or abnormally augment the occurrence-frequency of anomalous phenotypes outside of the natural variation in the species. Such practices cause rapid depletion of the normal genetic variation, the maintenance of which is a stated goal of the professional zoological community, while simultaneously subjecting some of the animals that are produced to, now documented, poor welfare, poor health, and reduced fitness.

Breeding practices that increase the phenotypic expression of single rare alleles through intentional inbreeding cause abnormal or aberrant external and internal conditions and characteristics. The predictability of such outcomes from intentional inbreeding to produce phenotypic anomalies indicates that these practices are not in adherence to AZA's Board-approved Policy on the Presentation of Animals (22 July 2008). Thus, AZA-accredited institutions should not engage in intentional inbreeding practices for the purpose of producing anomalous phenotypes from the perspectives of welfare, education, population management, and conservation.

## References

- Ballou JD, Lacy RC. 1995. Identifying genetically important individuals for management of genetic diversity in pedigreed populations. In J.D. Ballou, M. Gilpin, and T.J. Foose (eds.), Population Management for Survival & Recovery. Analytical Methods and Strategies in Small Population Conservation. Columbia University Press, New York, NY: pp. 76-111.
- Brandl P. 2010. Generically speaking. ZOOQUARIA (EAZA news) Issue 70 (summer 2010):19-22.
- Creel DJ, Giolli RA. 1972. The primary optic projection in pigmented and albino guinea pigs: an experimental degeneration study. Exp Neurol 30:25-39.
- Council of Europe: European Convention for the protection of pet animals, 13. Nov. 1987 (ETS 125), Council of Europe, Strasbourg-Cedex, 1987 ([www.bvet.ch](http://www.bvet.ch), [www.coe.int](http://www.coe.int)).
- Council of Europe: Resolution on the breeding of pet animals, Multilateral Consultation of parties to the European Convention for the protection of pet animals (ETS 125), March 1995 in Strasbourg, Document CONS 125(95)29, Council of Europe, Strasbourg-Cedex, 1995b ([www.bvet.ch](http://www.bvet.ch)).
- Council of Europe: Declaration of intent, Multilateral Consultation of parties to the European Convention to the European Convention for the protection of pet animals (ETS 125), March 1995 in Strasbourg, Document CONS 125(95)29, Council of Europe, Strasbourg-Cedex, 1995a ([www.bvet.ch](http://www.bvet.ch)).
- Falconer DS. 1981. Introduction to quantitative genetics. Second edition. London, New York: Longman: p.340.
- Federation of Veterinarians of Europe (FVE): Resolution "Breeding and Animal Welfare", FVE/99/010, 1999 ([www.fve.org](http://www.fve.org)).
- Fox PR. 1988. Congenital feline heart disease. In: Canine and Feline Cardiology. Ed. Fox PR. Churchill, Livingstone Inc., New York, NY: pp. 404-8.
- Goldin JP, Lambrecht NE. 1999. Double aortic arch and persistent left vena cava in a white lion cub (*Panthera leo*). J Zool Wildl Med 30:145-150.
- Guillery RW. 1986. Neural abnormalities of albinos. Trends Neurosci 9:364-367.
- Guillery RW, Kaas JH. 1973. Genetic abnormality of the visual pathways in a "white tiger". Science 180:1287-1288.
- Ketz CJ, Radlinsky M, Armbrust L, Carpenter JW, Isaza R. 2001. Persistent right aortic arch and aberrant subclavian artery in a white Bengal tiger (*Panthera tigris*). J of Zoo Wildl Med 32(2):268-272.
- Kirkwood JK, Weddell S, Hubrecht RC, Wickens SM. 2010. Darwinian selection, selective breeding and the welfare of animals. Proceedings of the UFAW international symposium, Bristol 2009. Animal Welfare 19(S): pp.1-149.
- Kliot M, Shatz CJ. 1985. Abnormal development of the retinogeniculate projection in Siamese Cats. J of Neuroscience 5(10): 2641-2653.
- Lacy R. 2000. Should we select genetic alleles in our conservation breeding programs? Zoo Biol 19: 279-282.
- Laikre L. 1999. Hereditary defects and conservation management of captive populations. Zoo Biol 18:81-99.
- Latinen K. 1987. White tigers and species survival plans. In: Tigers of the world: The Biology, Biopolitics, Management and Conservation of an Endangered Species, eds. RL Tilson and US Seal, Noyes Publ., Park Ridge, NJ: pp.391-393.
- Leyhausen P, Reed TH. 1971. The white tiger: care and breeding of a genetic freak. Smithson 2:24-31.
- Not I. 1998. Beurteilung verschiedener Zuchtlinien in Ziervögeln, Kleinnagern, Zierfischen und Reptilien in tierschützerischer Hinsicht. Dissertation (PhD Thesis). University of Zürich.



- Not I, Isenbügel E, Bartels T, Steiger A. 2008. Assessment of animal welfare aspects in extreme breeds of pet animals: small mammals, ornamental birds, fish, reptiles. (article in German). Schweiz Arch Tierheilk 150(5):235-241.
- Rooney NJ, Sargan DR. 2010. Welfare concerns associated with pedigree dog breeding in the UK. Animal Welfare 19(S):133-140.
- Roychoudhury AK. 1978. A study of inbreeding in white tigers. Sci Cult 44:371-2.
- Roychoudhury AK, Sankhala KS. 1979. Inbreeding in white tigers. Proc Indian Acad Sci 88B, Part I, No. 5:311-323.
- Sanderson KJ, Guillery RW, and RM Shackelford. 1974. Congenitally abnormal pathways in mink with reduced retinal pigment. J Comp Neurol 154:225-245.
- Schroeter W. 1981. On color, color deviation, stripe diminuation and color-brightening in the tiger, *Panthera tigris*. Säugetierkunde Mitt 29(4):1-8.
- Steiger A. 2005. Breeding and welfare of cats. In: The welfare of cats. Ed. I. Rochlitz, Springer: pp. 259-276.
- Steiger A. 2008. Assessment of animal welfare aspects in extreme breeds of pet animals: Principles, rules and other measures. (article in German). Schweiz Arch Tierheilk 150 (5):211-216.
- Steiger A, Stucki F, Peyer N, Keller P. 2008. Assessment of animal welfare aspects in extreme breeds of pet animals: dogs and cats. (article in German). Schweiz Arch Tierheilk 150(5):217-225.
- Stucki F, Bartels T, Steiger A. 2008. Assessment of animal welfare aspects in extreme breeds of rabbits, poultry and pigeons. (article in German). Schweiz Arch Tierheilk 150(5):227-234.
- Thornton IWB, Young KK, Sankhala KS. 1967. The genetics of the white tigers of Rewa. J of Zool 152:127-135.
- Thornton IWB. 1978. White tiger genetics – further evidence. J Zool 185(3):389-94.