


Chemists achieve molecular first

April 15 2014



This schematic shows how the three molecules are interlocked through a single point to produce a [3]catenane.

(Phys.org) —Chemists from Trinity College Dublin have achieved a long-pursued molecular first by interlocking three molecules through a single point. Developing interlocked molecules is one of the greatest challenges facing researchers, and the Trinity chemists' achievement represents the first time three molecules have been linked in a non-linear 'chain-like' form.

Interlocked molecules have major applications in nanoscience, as they can be used as molecular shuttles and switches, and because they can function as molecular motors, mimicking the action of many biological

systems.

Molecules that are interlocked together are unique in that they are not connected by any chemical bonds, which give other compounds their individual, defined structures following chemical reactions. Instead, the interlocked molecules typically exist as rings that together form a chain, like the pattern seen on the front cover of the iconic Book of Kells.

Led by Professor of Chemistry at Trinity, Thorfinnur Gunnlaugsson, the work was carried out by PhD student, Dr Christophe Lincheneau, who is now a postdoctoral fellow at CEA Grenoble. Dr Lincheneau used a metal 'lanthanide' ion called Europium and a catalyst developed by the Nobel Laureate, Professor Robert Grubbs of Caltech, to interlock three molecules through a single point. The important discovery was recently published in the high-impact journal of the Royal Society of Chemistry, *Chemical Communications*. It was also featured in *Nature Chemistry's* April issue as one of three Research Highlights.

Professor Gunnlaugsson said: "This work opens up a new avenue for developing complex supramolecular self-assembly structures. The luminescent properties of the lanthanide were very important to our study, as they allowed us to monitor the self-assembly processes between Europium and the molecules in real-time."

Europium luminescence is a powerful tool with wider applications. For example, it is currently being used to aid authorities in the prevention of counterfeiting, while luminescent lanthanide ion complexes are employed in the various denominations of Euro notes as red and green-emitting dyes.

Professor Gunnlaugsson's research group, which is located in the Trinity Biomedical Sciences Institute, was also able to accurately determine the mass of their interlocked '[3]catenane' by using the School of

Chemistry's state-of-the-art nuclear magnetic resonance (NMR) spectroscope and mass spectrometry facilities, which are part-funded under the HEA PRTL I Programmes. Their published research was funded by Science Foundation Ireland under the 2010 Principle Investigation Programme.

"We are now actively pursuing the development of other interlocked [molecules](#) and self-assembly structures using the lanthanide template design strategy we have developed and discussed in our recent publication. We hope this is just the first of many exciting and important discoveries," added Professor Gunnlaugsson.

More information: "Self-assembly formation of mechanically interlocked [2]- and [3]catenanes using lanthanide ion [Eu(III)] templation and ring closing metathesis reactions." Christophe Lincheneau, Bernard Jean-Denis, Thorfinnur Gunnlaugsson. *Chem. Commun.*, 2014,50, 2857-2860. [DOI: 10.1039/C3CC49640F](#). Received 19 Dec 2013, Accepted 22 Jan 2014. First published online 23 Jan 2014

Provided by Trinity College Dublin

Citation: Chemists achieve molecular first (2014, April 15) retrieved 4 October 2025 from <https://phys.org/news/2014-04-chemists-molecular.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
