


Researchers develop model to study heavy-quark recombination in quark-gluon plasma

July 11 2024



Quark recombination in the quark-gluon plasma formed in high-energy nuclei collisions enhances the production of B_c mesons. These mesons consist of a charm quark and a bottom quark. Most of the quark-gluon plasma decays into thousands of other particles. Credit: CERN (CMS Collaboration, B. Wu, Z. Tang, M. He, and R. Rapp)

Researchers from the HEFTY Topical Collaboration investigated the recombination of charm and bottom quarks into B_c mesons in the quark-gluon plasma (QGP). They have developed a transport model that simulates the kinetics of heavy-quark bound states through the expanding QGP fireball formed in high-energy heavy-ion collisions. Previous research has successfully used this model to describe the production of charm-anticharm and bottom-antibottom bound states, and thus can provide predictions for B_c particles (charm-antibottom bound states).

The paper is [published](#) in the journal *Physical Review C*.

A QGP formed in high-energy heavy-ion collisions only lasts for a short time before disintegrating into thousands of particles that can be observed in detectors. These detectors track signatures—the signals produced by specific types of particles. The discovery and study of QGP formation in heavy-ion experiments requires signatures that do not occur in other types of collisions, such as [proton-proton collisions](#).

In this study, researchers carried out theoretical simulations of charm and bottom quarks diffusing through the QGP. They found that the recombination of these quarks enhances the production of B_c mesons. This mechanism does not occur in proton-proton collisions and thus can serve as a clean signature of QGP formation.

The researchers used realistic spectra of charm and bottom quarks, computed from their diffusion through the QGP, to evaluate their recombination processes. The results show a large enhancement of the B_c yield in collisions of lead (Pb) nuclei, relative to that in proton collisions. The largest effect is predicted for slow-moving B_c mesons in "head-on" collisions of the Pb nuclei, where a large QGP fireball with appreciable numbers of charm and bottom quarks is formed.

The [theoretical calculations](#) agree with pioneering data of the CMS collaboration at the Large Hadron Collider (LHC). However, the data are not yet sensitive to slow-moving B_c mesons; future data will therefore provide a critical test of this QGP signature.

More information: Biaogang Wu et al, Recombination of B_c mesons in ultrarelativistic heavy-ion collisions, *Physical Review C* (2024). [DOI: 10.1103/PhysRevC.109.014906](#)

Provided by US Department of Energy

Citation: Researchers develop model to study heavy-quark recombination in quark-gluon plasma (2024, July 11) retrieved 4 October 2025 from <https://phys.org/news/2024-07-heavy-quark-recombination-gluon-plasma.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
--